

Environmental Protection Agency

§ 180.127

hulls as animal feed. Peanut hulls containing residues of inorganic bromides from the use of methyl bromide are unsuitable as an ingredient in the feed of meat and dairy animals and should not be represented, sold, or used for that purpose.

[58 FR 65555, Dec. 15, 1993]

§ 180.124 Methyl bromide; tolerances for residues.

(a) *General.* A tolerance is established for residues of the fumigant methyl bromide, including metabolites and degradates, in or on the commodity in the table below. Compliance with the tolerance level specified below is to be determined by measuring only methyl bromide.

Commodity	Parts per million
Cotton, undelinted seed	150

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[77 FR 35298, June 13, 2012]

§ 180.127 Piperonyl butoxide; tolerances for residues.

(a) *General.* (1) Tolerances for residues of the insecticide piperonyl butoxide [(butyl carbityl)(6-propyl piperonyl)ether] are established in or on the following food commodities:

Commodity	Parts per million
Almond, postharvest	8
Apple, postharvest	8
Barley, postharvest	20
Bean, postharvest	8
Birdseed, mixtures, postharvest	20
Blackberry, postharvest	8
Blueberry, postharvest	8
Boysenberry, postharvest	8
Buckwheat, grain, postharvest	20
Cattle, fat	0.1
Cattle, meat	0.1
Cattle, meat byproducts	0.1
Cherry, sweet, postharvest	8
Cherry, tart, postharvest	8
Cocoa bean, roasted bean, postharvest	8
Coconut, copra, postharvest	8
Corn, field, grain, postharvest	20
Corn, pop, postharvest	20
Cotton, undelinted seed, postharvest	8
Crabapple, postharvest	8
Currant, postharvest	8
Dewberry, postharvest	8

Commodity	Parts per million
Egg	1
Fig, postharvest	8
Flax, seed, postharvest	8
Goat, fat	0.1
Goat, meat	0.1
Goat, meat byproducts	0.1
Gooseberry, postharvest	8
Grape, postharvest	8
Guava, postharvest	8
Hog, fat	0.1
Hog, meat	0.1
Hog, meat byproducts	0.1
Horse, fat	0.1
Horse, meat	0.1
Horse, meat byproducts	0.1
Loganberry, postharvest	8
Mango, postharvest	8
Milk, fat	0.25
Muskmelon, postharvest	8
Oat, postharvest	8
Orange, postharvest	8
Peach, postharvest	8
Peanut, postharvest	8
Pea, postharvest	8
Pear, postharvest	8
Pineapple, postharvest	8
Plum, prune, fresh, postharvest	8
Potato, postharvest	0.25
Poultry, fat	3
Poultry, meat	3
Poultry, meat byproducts	3
Raspberry, postharvest	8
Rice, postharvest	20
Rye, postharvest	20
Sheep, fat	0.1
Sheep, meat	0.1
Sheep, meat byproducts	0.1
Sorghum, grain, postharvest	8
Sweet potato, postharvest	0.25
Tomato, postharvest	8
Walnut, postharvest	8
Wheat, postharvest	20

(2) Piperonyl butoxide may be safely used in accordance with the following prescribed conditions:

(i) It is used or intended for use in combination with pyrethrins for control of insects:

(A) In cereal grain mills and in storage areas for milled cereal grain products, whereby the amount of piperonyl butoxide is at least equal to but not more than 10 times the amount of pyrethrins in the formulation.

(B) On the outer ply of multiwall paper bags of 50 pounds or more capacity in amounts not exceeding 60 milligrams per square foot, whereby the amount of piperonyl butoxide is equal to 10 times the amount of pyrethrins in the formulation. Such treated bags are to be used only for food, dried.

(C) On cotton bags of 50 pounds or more capacity in amounts not exceeding 55 milligrams per square foot of